



KENVET AFRI LIMITED

Products Catalogue



P.o. Box 907-00606, Sarit Centre, Nairobi, Kenya.

Table of Contents

KenVaq.....	1
KenVasone.....	1
Tetraken 5%.....	2
Teraken 10%.....	2
Tetraken 20%.....	3
KenVasole 7.5%.....	3
Kenroxin 10%.....	4
KenBoost.....	5
KenFer.....	5
KenStrep-P.....	6
KenStrep-LA.....	6
KenVectin.....	7
Tyloken.....	7
Kenbezole 300mg.....	8
Kenbezole 600mg.....	8
Kenbezole 2500mg.....	9
Pippraz.....	9
Kenbezole 2.5%.....	10
Kenbezole 10%.....	10
KenDox-T.....	11
Saltrim Vet.....	11
Amproken.....	12
Tetraken Chick Formula.....	12
Tetraken Egg Formula.....	13
Tetraken Soluble Powder.....	13
KenBoost WS.....	14
KenBoost Extra.....	14
Livtrim Bolus.....	15
Mebpip-P.....	15
Kenmidine.....	16
KenVasole.....	16
KenOverm DS.....	17
KenTraz.....	17
Samovet.....	18
Tetraken Aerosol.....	18
KenVaq Plus.....	19



KENVET AFRI LIMITED

P.o. Box 907-00606, Sarit Centre, Nairobi, Kenya.

KenVaq



Composition

Each ml contains
Buparvaquone 50mg

Indications

For the treatment of East Coast Fever and other forms of Theileriosis in cattle caused by strains of *Theileria parva*, *Theileria annulata*, *Theileria mutans* and *Theileria orientalis*.

Dosage & Administration

A single injection of 1ml KenVaq per 20kg bodyweight (2.5mg buparvaquone per kg body weight) by deep intramuscular injection into the neck muscles.

Precautions:

Do not administer by intravenous or subcutaneous route.

Withdrawal Period from the last day of treatment:

Meat: 42 days | Milk: 48 hours

Storage

Store in a dry place, below 30°C, protected from direct sunlight.
Keep all medicines out of reach of children.

Presentation

50ml & 100ml

FOR VETERINARY USE ONLY

KenVasone



Composition

Each 1ml contains:
Dexamethasone sodium phosphate equivalent to 2mg dexamethasone
Excipients q.s.

Indications

Metabolic disorders, non-infectious inflammatory processes, especially acute musculoskeletal inflammations, allergic conditions, stress and shock conditions.

As an aid in infectious diseases.

Dosage & Administration

For intravenous or intramuscular injection.
Normal aseptic precautions should be observed.

Cattle, Horses: 1ml per 25kg body weight.
Calves, Dogs: 1ml per 10kg body weight.
e.g.

Cattle:	400Kg	16ml
Horses:	500Kg	20ml
Cats:	5Kg	0.5ml
Dogs:	10Kg	1ml

Withdrawal periods from the last day of treatment:

Meat: 21 days | Milk: 72 hours

Storage

Store in a dry place, below 30°C, protected from direct sunlight.
Keep all medicines out of reach of children.

Presentation

50ml & 100ml

FOR VETERINARY USE ONLY



KENVET AFRI LIMITED

P.o. Box 907-00606, Sarit Centre, Nairobi, Kenya.

Tetraken 5%



Composition

Each ml contains:
Oxytetracycline 50mg (as Hydrochloride) BP Vet.

Indications

A broad-Spectrum Antimicrobial agent against Gram-Positive and Gram-Negative bacteria, protozoa and rickettsia.

Dosage & Administration

For intramuscular, subcutaneous or slow intravenous injection.

Horses, Cattle:

1ml per 5-10kg body weight daily for 3-5 days.

Pigs, Sheep, Goats:

5-8ml per 50kg body weight daily for 3-5 days.

Dogs, Cats:

1ml per 5kg body weight daily for 3-5 days.

Withdrawal period from the last day of treatment:

Milk: 3 Days

Meat: 20 Days

Use the product within 28 days after withdrawal of first dose.

Storage

Store in a dry place, below 30°C, protected from direct sunlight.

Keep all medicines out of reach of children.

Presentation

50ml & 100ml

FOR VETERINARY USE ONLY

Tetraken 10%



Composition

Each ml contains:
Oxytetracycline 100mg (as Hydrochloride) BP Vet.

Indications

A broad-Spectrum Antimicrobial agent against Gram-Positive and Gram-Negative bacteria, protozoa and rickettsia.

Dosage & Administration

For intramuscular, subcutaneous or slow intravenous injection.

Horses, Cattle:

1ml per 5-10kg body weight daily for 3-5 days.

Pigs, Sheep, Goats:

5-8ml per 50kg body weight daily for 3-5 days.

Dogs, Cats:

1ml per 5kg body weight daily for 3-5 days.

Withdrawal period from the last day of treatment:

Milk: 3 Days

Meat: 20 Days

Use the product within 28 days after withdrawal of first dose.

Storage

Store in a dry place, below 30°C, protected from direct sunlight.

Keep all medicines out of reach of children.

Presentation

50ml, 100ml & 250ml

FOR VETERINARY USE ONLY



KENVET AFRI LIMITED

P.o. Box 907-00606, Sarit Centre, Nairobi, Kenya.

Tetraken 20%



Composition

Each ml contains:
Oxytetracycline 200mg (as Hydrochloride) BP Vet.

Indications

A broad-Spectrum Antimicrobial agent against Gram-Positive and Gram-Negative bacteria, protozoa and rickettsia.

Dosage & Administration

For intramuscular, subcutaneous or slow intravenous injection.

Horses, Cattle:

1ml per 10kg body weight by deep intramuscular injection.

Withdrawal period from the last day of treatment:

Milk: 3 Days

Meat: 28 Days

Use the product within 28 days after withdrawal of first dose.

Storage

Store in a dry place, below 30°C, protected from direct sunlight.

Keep all medicines out of reach of children.

Presentation

50ml, 100ml & 250ml

FOR VETERINARY USE ONLY

KenVasole 7.5%



Composition

Levamisole Hydrochloride 75mg
Excipients up to 1ml

Indications

For treatment and control of nematode infections.

Dosage & Administration

By subcutaneous injection.

1ml per 10kg body weight.

Cattle: 10ml per 100kg body weight (max. 20ml per injection site).

Sheep: 1ml per 10kg body weight.

Withdrawal period from the last day of treatment:

Not to be used in animals producing milk for human consumption.

Meat: Cattle 28 days

Sheep: 15 days.

Caution:

Use the product within 28 days after withdrawal of first dose.

Storage

Store in a dry place, below 30°C, protected from direct sunlight.

Keep all medicines out of reach of children.

Presentation

50ml & 100ml

FOR VETERINARY USE ONLY



KENVET AFRI LIMITED

P.o. Box 907-00606, Sarit Centre, Nairobi, Kenya.

Kenroxin 10%



Composition

Each ml contains:
Enrofloxacin 100mg
Excipients qs to 1ml

Indications

Kenroxin 10% injection is indicated against gastrointestinal, respiratory and reproductive infections caused by micro-organisms susceptible to enrofloxacin, like broncho-pneumonia, colisepticemia/coli diarrhoea, mycoplasma infections, enteritis, metritis, mastitis, pasteurellosis, infections of the urinary tract, wound infections, paratyphoid and secondary bacterial infections such as complications of viral diseases in cattle, sheep, goats and pigs.

Dosage & Administration

Kenroxin 10% injection may be administered as a single dose (cattle & swine) or for multiple days (cattle) of therapy. Selection of the appropriate dose and duration of therapy should be based on assessment of the severity of the disease, clinical response and pathogen susceptibility.

Single Dose Therapy (Cattle): Administer once 7.5-12.5mg/kg or 7.5-12.5ml/100kg body weight subcutaneously.

Multiple Day Therapy (Cattle): Administer daily 2.5-5mg/kg or 2.5-5ml/100kg body weight at 24 hours interval for 3 days. Administered dose volume should not exceed 20ml per injection site.

Sheep & Goats:

Administer once 7.5mg/kg or 7.5ml/100kg body weight subcutaneously. Administered dose volume should not exceed 5ml per injection site.

Swine:

Administered once behind the ear a subcutaneous dose of 7.5mg/kg or 7.5ml/100kg body weight. Administered volume should not exceed 5ml per injection site.

Precautions:

To be used by or on order of a licensed veterinarian. Keep all medicines out of reach of children

Withdrawal period from the last day of treatment:

Cattle: 28 days.
Swine, Goats and Sheep: 5 days

Storage

Store in a dry place, below 30°C, protected from direct sunlight. Keep all medicines out of reach of children.

Presentation

50ml & 100ml

FOR VETERINARY USE ONLY



KENVET AFRI LIMITED

P.o. Box 907-00606, Sarit Centre, Nairobi, Kenya.

KenBoost



Composition

Vitamin A 3,000 IU, Vitamin D3 2,000 IU, Vitamin E 4mg, Vitamin B12 10mcg, Ascorbic acid 1mg, Vitamin B1 10mg, Vitamin B6 5mg, Vitamin B2 1mg, Nicotinamide 12mg, Pantothenal 10mg and D-Biotin 10mcg

Indications

Prevention and treatment of vitamin deficiencies in horses, cattle, sheep & goats, particularly during periods of illness, convalescence and general unthriftiness.

Dosage & Administration

By intramuscular or subcutaneous injection. Clean the area of the injection site and swab with spirit. Ensure that all injection equipment is thoroughly clean and sterilised before use. Large Animals: 2ml per 50Kg
Small Animals: 1ml per 10Kg
The period of time between the withdrawal of the first and last dose from the container should not be unduly prolonged.
Withdrawal period from last day of treatment
Milk: Nill | Meat: Nill

Storage

Store in a dry place, below 30°C, protected from direct sunlight.
Keep all medicines out of reach of children.

Presentation

50ml & 100ml

FOR VETERINARY USE ONLY

KenFer



Composition

Each ml contains:
Iron Dextran equivalent to 100 mg Iron

Indications

Prevention and treatment of iron deficiency in Piglets and other animals.
It is contraindicated in cases of vitamin E or Selenium deficiency in Piglets.

Dosage & Administration

By deep intramuscular injection
Piglets (prevention): 2ml on 3 day of life and repeat dose on 21st Day.
Cattle (Treatment): 4-8ml
Calves (Treatment): 3-6ml

Withdrawal period from the last day of treatment:
None

Storage

Store in a dry place, below 30°C, protected from direct sunlight.
Keep all medicines out of reach of children.

Presentation

50ml & 100ml

FOR VETERINARY USE ONLY



KENVET AFRI LIMITED

P.o. Box 907-00606, Sarit Centre, Nairobi, Kenya.

KenStrep-P



Composition

Each ml of suspension for injection contains:
200mg Procaine Benzylpenicillin and 200mg Dihydrostreptomycin (as sulphate).

Indications

For the treatment of infections caused by bacteria sensitive to penicillin and dihydrostreptomycin.

Dosage & Administration

For intramuscular administration only.
Cattle: 1 ml per 20 kg body weight for 3 days.
Calves, goats, sheep and
Swine: 1 ml per 10 kg body weight for 3 days.
Note: Shake well before use and do not administer more than 20ml in cattle, more than 10ml in swine and more than 5ml in calves, sheep and goats per injection site.
Clean the area of injection and swab with spirit.
Withdrawal period from the last day of treatment:

For kidney: 45 days

For meat: 21 days

For milk: 3 days

The withdrawal period for milk relates to cattle only.

Storage

Store in a dry place, below 30°C, protected from direct sunlight.

Keep all medicines out of reach of children.

Presentation

50ml & 100ml

FOR VETERINARY USE ONLY

KenStrep LA



Composition

Contains per ml:

Procaine Penicillin G.....100 000 IU

Benzathine Penicillin G.....100 000 IU

Dihydrostreptomycin Sulphate.....200mg

Excipients ad.....1ml

Indications

Arthritis, mastitis and gastrointestinal, respiratory and urinary tract infections caused by penicillin and dihydrostreptomycin sensitive micro-organisms like campylobacter, clostridium, carynebacterium, E coli, erysipelothrix, Haemophilus, Klebsiella, Listeria, Pasteurella, Salmonella, Staphylococcus and streptococcus spp.

Dosage & Administration

For intramuscular administration.

General: 1ml per 10kg body weight every 72 hours.

When recommended by professional opinions, it can be administered every 48 hours.

Withdrawal period from the last day of treatment: For Kidney: 45 days

For Meat: 30 days

For Milk: 5 days

Storage

Store in a dry place, below 30°C, protected from direct sunlight.

Keep all medicines out of reach of children.

Presentation

50ml & 100ml

FOR VETERINARY USE ONLY



KENVET AFRI LIMITED

P.o. Box 907-00606, Sarit Centre, Nairobi, Kenya.



Composition

Each ml contains: Ivermectin 10 mg

Indications

KenVectin provides treatment and control of Gastrointestinal Worms, Lung Worms, Eye Worms, Warbles, Mange Mites and Sucking Lice.

Dosage & Administration

For subcutaneous administration only.
Cattle, Sheep, Goat and Camels: 1.0ml per 50kg body weight.
Swine:
1.0ml per 33kg body weight.
Use the product within 28 days after withdrawal of first dose.

Withdrawal period from the last day of treatment:

Meat: Cattle: 28 Days
Pigs: 21 Days
Goats and Sheep: 21 Days

Storage

Store in a dry place, below 30°C, protected from direct sunlight.
Keep all medicines out of reach of children.

Presentation

10ml, 50ml & 100ml
FOR VETERINARY USE ONLY



Composition

Each ml contains:
Tylosin Tartrate 200mg

Indications

Infections caused by micro-organisms sensitive to Tyloken, including Respiratory infections, Vibronic Dysentery in Pigs, Mycoplasmosis, Foot rot, Mastitis, Metritis, Cystitis, Post-operative infections in Cattle, Pigs, Sheep Goats.

Dosage & Administration

For intramuscular or subcutaneous injection.
Cattle: 3.0-5.0ml per 100kg body Weight daily.
Calf: 1.5-2.0ml per 50kg body Weight daily.
Pig: 1.2-2.5ml per 50kg body weight.
All doses for 1-5 consecutive days.

Withdrawal periods from last day of treatment:

Milk: 60 hrs
Meat: Cattle 23 days Pigs: 18 days Sheep: 31 days

Storage

Store in a dry place, below 30°C, protected from direct sunlight.
Keep all medicines out of reach of children.

Presentation

50ml & 100ml
FOR VETERINARY USE ONLY



Kenbezole Bolus 300mg



Composition

Each uncoated bolus contains
Albendazole 300mg.

Indications

For effective control and treatment against gastrointestinal roundworms, lungworms, liverflukes and tapeworms infestation in cattle, camels, goats and sheep.

Dosage & Administration

Administer orally.

Body Weight	Worms 7.5mg/Kg	Liver Flukes 15mg/kg
20 Kg	0.5 Bolus	1 Bolus
40 Kg	1 Bolus	2 Bolus
60 Kg	1.5 Bolus	3 Bolus

Withdrawal Periods: Meat: 12 days, Milk: 4 days.

Caution

Do not treat animals during first month of pregnancy. Do not under dose.

Storage

Store in a dry place, below 30°C, protected from direct sunlight.

Keep all medicines out of reach of children.

Presentation

10 x 5 boli

FOR VETERINARY USE ONLY

Kenbezole Bolus 600mg



Composition

Each uncoated bolus contains
Albendazole 600mg.

Indications

For effective control and treatment against gastrointestinal roundworms, lungworms, liverflukes and tapeworms infestation in cattle, camels, goats and sheep.

Dosage & Administration

Administer orally.

Body Weight	Worms 7.5mg/Kg	Liver Flukes 15mg/kg
120 Kg	1.5 Bolus	3 Boli
160 Kg	2 Boli	4 Boli
200 Kg	2.5 Boli	5 Boli
240 Kg	3 Boli	6 Boli

Withdrawal Periods: Meat: 12 days, Milk: 4 days.

Caution

Do not treat animals during first month of pregnancy. Do not under dose.

Storage

Store in a dry place, below 30°C, protected from direct sunlight.

Keep all medicines out of reach of children.

Presentation

10 x 5 boli

FOR VETERINARY USE ONLY



KENVET AFRI LIMITED

P.o. Box 907-00606, Sarit Centre, Nairobi, Kenya.

Kenbezole Bolus 2500mg



Composition

Each uncoated bolus contains
Albendazole 2500mg.

Indications

For effective control and treatment against gastrointestinal roundworms, lungworms, liverflukes and tapeworms infestation in cattle, camels, goats and sheep.

Dosage & Administration

Administer orally.

Body Weight	Worms 7.5mg/Kg	Liver Flukes 15mg/kg
165 Kg	0.5 Bolus	1 Bolus
330 Kg	1 Bolus	2 Boli
500 Kg	1.5 Bolus	3 Boli

Withdrawal Periods: Meat: 12 days, Milk: 4 days.

Caution

Do not treat animals during first month of pregnancy. Do not under dose.

Storage

Store in a dry place, below 30°C, protected from direct sunlight.


Keep all medicines out of reach of children.

Presentation

10 x 5 boli

FOR VETERINARY USE ONLY

Pippraz



Composition

Each gram contains:
Piperazine Citrate 360mg.

Indications

For Prompt Expulsion of Roundworms & Pinworms in Cattle, Horses, Pigs & Poultry

Dosage & Administration

Directions for use

Administration:

Give as a 1 day treatment.

The preparation for poultry can be kept for 2 days.

General dosage instructions:

Pigs, Horses and Cattle: 1-2 gm per 10 kg body weight.

Poultry:

Dissolve 30 gm in sufficient quantity of water (20-30) litres per 100 birds or (1.5 level teaspoonfuls) in 3-5 litres of water per 15 birds.

Storage

Store in a dry place, below 30°C, protected from direct sunlight.

Keep all medicines out of reach of children.

Presentation

100gm


FOR VETERINARY USE ONLY



KENVET AFRI LIMITED

P.o. Box 907-00606, Sarit Centre, Nairobi, Kenya.

KenBezole 2.5%



Composition

Each ml Contains:
Albendazole 25mg

Indications

For prevention & treatment of roundworm, tapeworm and liver fluke infestations in cattle, sheep and goats.

Dosage & Administration

Kenbezole 2.5%w/v Suspension is given by oral route using a drenching gun or a dosing syringe.

Withdrawal periods from the last day of treatment:

Meat: 14 days

Milk: 3 days

Storage

Store in a dry place, below 30°C, protected from direct sunlight.


Keep all medicines out of reach of children.

Presentation

120ml, 500ml, 1L & 5L

FOR VETERINARY USE ONLY

KenBezole 10%



Composition

Each ml Contains:
Albendazole 100mg

Indications

For prevention & treatment of roundworm, tapeworm and liver fluke infestations in cattle, sheep and goats.

Dosage & Administration

Kenbezole 10%w/v Suspension is given by oral route using a drenching gun or a dosing syringe.

Withdrawal periods from the last day of treatment:

Meat: 14 days

Milk: 3 days

Storage

Store in a dry place, below 30°C, protected from direct sunlight.

Keep all medicines out of reach of children.

Presentation

120ml, 500ml, 1L & 5L

FOR VETERINARY USE ONLY



KENVET AFRI LIMITED

P.o. Box 907-00606, Sarit Centre, Nairobi, Kenya.



Composition

Each 1 gram contains:
 Doxycycline Hyclate.....200mg.
 Tylosin Tartrate.....200mg.

Indications

Respiratory and gastrointestinal infections caused by Tylosin and Doxycycline sensitive organisms like Staphylococcus, Mycoplasma, E coli, Salmonella, Pasteurella, Bordetella, Streptococcus, Campylo- bacter and Haemophilus spp. in poultry, swine, sheep, goats and calves.

Dosage & Administration

For oral administration.
 Poultry: 2.5gm in 5 litres drinking water for 3-5 days.
 Pigs, Calves, Sheep & Goats: 2.5gm per 50kg body weight for 3-5 days.
 Note: For pre-ruminant calves, lambs and kids only.

Withdrawal Period From Last Day Of Treatment Meat:

Poultry: 7 days | Swine: 8 days
 Calves, Sheep & Goats: 14 days
 Eggs: 4 days

Storage

Store in a dry place, below 30°C, protected from direct sunlight.
 Keep all medicines out of reach of children.

Presentation

30gm & 100gm
 FOR VETERINARY USE ONLY



Composition

Trimethoprim BP 2.0%
 Sulphamethoxazole BP 10.0%

Indications

For the treatment of Infectious Coryza, Salmonella Infections, Respiratory Disease and Caecal Coccidiosis

Dosage & Administration

Add 5 gm Saltrim Vet to 5 litres of water and give to the birds for 5 days continuously or as directed by a veterinarian.

Precautions and Warnings

Do not give the birds any other drinking water over the treatment period. Give only the medicated water.
 Prepared medicated water to be used within 24 hours.

Withdrawal Periods

Meat: 6 days.
 Eggs: Should not be used in laying birds producing eggs for human consumption.

Storage


Store in a dry place, below 30°C, protected from direct sunlight.
 Keep all medicines out of reach of children.

Presentation

30gm & 100gm
 FOR VETERINARY USE ONLY



Amproken



Composition

Each gram contains:

- Amprolium Hydrochloride BP (Vet) 200mg,
- Sulfaquinoxaline (as Sodium) 150mg,
- Vitamin K3 5mg,
- Vitamin A, retinol-acetate 15,000 IU.

Indications

For Prevention & Treatment of Coccidiosis in Poultry & bacterial infections in Poultry, Calves, Sheep and Goats.

Dosage & Administration

For oral administration.

Calves, Sheep and Goats:

Preventive: 1 gram per 50-100kg body weight through drinking water or milk for 21 days.

Curative: 5 gram per 25-50kg body weight through drinking water or milk for 5 days.

Poultry: 20 gram per 20-40 litre drinking water or milk for 5 days.

Note: For pre-ruminant calves, lambs & kids only.

Do not administer to hens producing eggs for human consumption.

Withdrawal Periods from the the last day of treatment:

Meat: Calves, sheep, goats and poultry 14 days.

Storage

Store in a dry place, below 30°C, protected from direct sunlight.

Keep all medicines out of reach of children.

Presentation

30gm & 100gm

FOR VETERINARY USE ONLY

Tetraken Chick Formula



Composition

Each 25gm contains:

- Oxytetracycline HCL 1.375g, Vitamin A 55,000 I.U.
- Vitamin D3 10,000 I.U. Vitamin E 16.54mg,
- Vitamin K3 19.85mg, Vitamin B2 71.65mg,
- Vitamin B12 0.055mg, Nicotinamide 330.687mg,
- Calcium and Pantothenate 115.74mg.

Indications

Maintains appetite during stress periods like vaccination, extreme temperatures, change of feed and other upsets.

It also prevents infection and reduces mortality.

Dosage & Administration

For 100 starting chicks, use one 5ml spoonful per 5 litres drinking water or 2.5kg of feed daily for the first two weeks of life.

Withdrawal Period:

Eggs from treated animals should not be used for human consumption during treatment and 5 days after last day of treatment.

Storage

Store in a dry place, below 30°C, protected from direct sunlight.

Keep all medicines out of reach of children.

Presentation

25gm

FOR VETERINARY USE ONLY



KENVET AFRI LIMITED

P.o. Box 907-00606, Sarit Centre, Nairobi, Kenya.

Tetraken Egg Formula



Composition

Each 25gm contains:
Oxytetracycline HCL 1.375g, Vitamin A 55,000 I.U.,
Vitamin D3 10,000 I.U. Vitamin E 16.54mg,
Vitamin K3 19.85mg, Vitamin B2 71.65mg,
Vitamin B12 0.055mg, Nicotinamide 330.687mg,
Calcium and Pantothenate 115.74mg

Indications

For increasing egg production and protection against diseases.

Dosage & Administration

For young layers: Half 5ml teaspoonful in 5 litres of water or 2.5kg of feed at the start of laying for 5-7 days.

For older flocks: one 5ml teaspoonful in 5 litres of water or 2.5kg of feed at the start of laying for 5-7 days to maintain high egg production and combat diseases.

Withdrawal Period:

Eggs from treated animals should not be used for human consumption during treatment and 5 days after last day of treatment.

Storage

Store in a dry place, below 30°C, protected from direct sunlight.

Keep all medicines out of reach of children.

Presentation

25gm

FOR VETERINARY USE ONLY

Tetraken Soluble Powder



Composition

1g contains 50mg Oxytetracycline Hydrochloride.

Indications

Treatment of infections in livestock and poultry.

Dosage & Administration

Cattle, Calves, Fowls, Lambs, Sheep and Goats: 1 teaspoonful per 25kg body weight for 4-5 days.

Pigs: 2 teaspoonfuls in 5 litres of drinking water for 4-5 days:

Poultry: 1 to 2 teaspoonfuls in 5 litres of drinking water for 4-5 days.

Rabbits: 2 teaspoonfuls in 5 litres of drinking water for 4-5 day.

Prophylaxis: For disease prevention use half of the treatment dosage.

Note: For pre-ruminant calves, lambs and kids only.

Withdrawal Period from the last day of treatment:

Meat: 5 days

Milk: 1 day

Storage

Store in a dry place, below 30°C, protected from direct sunlight.

Keep all medicines out of reach of children.

Presentation

25gm

FOR VETERINARY USE ONLY



KENVET AFRI LIMITED

P.o. Box 907-00606, Sarit Centre, Nairobi, Kenya.

KenBoost WS



Composition

Each 30gm contains
Vitamin A 200,000I.U., Vitamin B1 60mg
Vitamin B2 60mg, Vitamin B6 60mg
Vitamin B12 600 mcg, Vitamin C 200mg
Vitamin D3 50,000I.U., Vitamin E 60I.U.
Vitamin K3 60mg, L (+) Choline Bitartrate
(Coated) 600mg, Pantothenic Acid 120mg
(as Calcium Pantothenate), Folic Acid 5mg
Niacinamide 420mg, Soluble Base to 30gm

Indications

- As a supplement in cases of vitamin deficiencies in all livestock.
- In cases of retarded growth.
- In times of convalescence.
- It may be used in combination with other drugs for control of diseases.
- To increase feed efficiency in all livestock.

Storage

Store in a dry place, below 30°C, protected from direct sunlight.
Keep all medicines out of reach of children.

Presentation

30gm & 100gm

FOR VETERINARY USE ONLY

KenBoost Extra



Composition

Each 30gm Powder Contains:
Vitamin A 457,500 IU, Vitamin D3 135,000 IU
Vitamin E 40.8 IU, Vitamin K 135 mg
Vitamin B2 135 mg, Vitamin B6 70.5 mg
Vitamin B12 345 mcg, Vitamin C 30mg
Niacin 502.5 mg, Methionine 306 mg
Lysine 457.5 mg, Zinc Sulphate 367.5 mg
Copper Sulphate 367.5 mg
Manganese Sulphate 367.5 mg
Sodium Chloride 1,500 mg
Sodium Sulphate 6,300mg
Potassium Chloride 2,640 mg
Soluble base (Dextrose) to 30 gm

Indications

- Provision of essential nutrients (vitamins, minerals & amino acids) in animals.
- Prevention of vitamin deficiencies and/or nutritional diseases in livestock especially young and debilitated animals.
- Improvement of egg production, weight gain, growth, production performance & reproduction in livestock.
- Management of stress resulting from various causes e.g. handling, transport, temperature fluctuation, vaccination etc.
- As an aid to recovery after treatment of coccidiosis, worm infestation, bacterial or viral infections.

Storage

Store in a dry place, below 30°C, protected from direct sunlight.
Keep all medicines out of reach of children.

Presentation

30gm & 100gm

FOR VETERINARY USE ONLY



KENVET AFRI LIMITED

P.o. Box 907-00606, Sarit Centre, Nairobi, Kenya.

Livtrim Bolus

Mebpip-P Worm Tablets



Composition

Each bolus contains:
Trimethoprim BP Vet 200mg and Sulfadiazine BP Vet 1g

Indications

Livtrim bolus is particularly effective against disease caused by E. Coli and Salmonella species of bacteria.

Livtrim bolus is used primarily for the oral treatment of bacterial scours in calves but may also be used for sheep, goats and foals. Livtrim bolus may also be used for systemic infections especially calf diphtheria, salmonellosis and bacterial pneumonia. Livtrim bolus may be used as pessaries for prevention and treatment of post-parturient infections including metritis and vaginitis in mares, cows, ewes and sows.

Dosage & Administration

For oral use.

Give Livtrim bolus by hand or balling gun or crush and disperse in water and administer as a drench.

Storage

Store in a dry place, below 30°C, protected from direct sunlight.

Keep all medicines out of reach of children.

Presentation

40 Boli

FOR VETERINARY USE ONLY



Composition

Each tablet contains:
Mebendazole BP 110mg
Piperazine Citrate BP 275mg
Praziquantel BP 25mg

Indications

Treatment for Roundworms, Hookworms and Tapeworms in dogs and cats.

Dosage & Administration

Dogs & Cats : One tablet per 5kg body weight. (Maximum 6 tablets for an adult dog).

- Tablets must be repeated 3-4 weeks after the first dose.
- Tablets are given by mouth directly or crushed and mixed with little amount of soup, milk or food.
- It is advisable to withhold food for 12 hours before administration of MEBPIP-P to get best results.
- As it is unlikely that unweaned animals become infected with tapeworms, MEBPIP-P is not recommended for unweaned puppies and kittens.

Storage

Store in a dry place, below 30°C, protected from direct sunlight.

Keep all medicines out of reach of children.

Presentation

6 Tablets

FOR VETERINARY USE ONLY



KENVET AFRI LIMITED

P.o. Box 907-00606, Sarit Centre, Nairobi, Kenya.

KenMidine



Composition

Contains:
33% w/v Sulfadimidine Sodium BP.

Indications

Coccidiosis; Helps in the control of outbreaks in chicken and turkeys.

Fowl Cholera, Coryza; Reduces the severity of the diseases and death losses in chicken and turkeys.

Fowl Typhoid; Reduces losses due to Salmonella gallinarum infections in poultry.

Dosage & Administration

125ml serves 400 birds per day.
Coccidiosis, Coryza, Fowl Cholera and Typhoid; Use 2 tablespoonfuls (20ml) in 5 litres of drinking water for 3-5 days.
Acute Cholera and Typhoid;
Use 2 tablespoonfuls (20ml) in 5 litres of drinking water for 6 consecutive days.

Caution:
With-holding period:
Eggs and Meat -96 hours after last treatment.

Storage

Store in a dry place, below 30°C, protected from direct sunlight.
Keep all medicines out of reach of children.

Presentation

125ml

FOR VETERINARY USE ONLY

KenVasole



Composition

Each 1ml contains
Levamisole HCL BP 15mg
Excipients qs

Indications

Kenvasole is a broad spectrum anthelmintic used in control and treatment of gastro-intestinal worms and lungworms infections in cattle, calves, sheep and goats.

Dosage & Administration

Dosage and Administration:
For oral administration only.
All species: 5ml per 10kg bodyweight.
It is safe to use in pregnant and lactating animals. Can be use during vaccinations and treatment of other conditions simultaneously. At the normal Kenvasole dose levels, animals rarely show any effects.
Occasionally some animals may show slight scouring. The effects may be enhanced in animals with severe liver damage.

Withdrawal Period From the last day of treatment:

Meat: 14 days | Milk: 1 day

Storage

Store in a dry place, below 30°C, protected from direct sunlight.
Keep all medicines out of reach of children.

Presentation

125ml, 500ml & 1L

FOR VETERINARY USE ONLY



KENVET AFRI LIMITED

P.o. Box 907-00606, Sarit Centre, Nairobi, Kenya.

KenOverm DS



Composition

Contains:

Oxyclozanide BP Vet 6.0% w/v, Levamisole Hydrochloride BP Vet 3.0% w/v and Cobalt Sulphate 0.764% w/v.

Indications

Kenoverm DS is a broad spectrum cattle and sheep anthelmintic indicated for the treatment and control of infections of both gastrointestinal and pulmonary nematodes (roundworms and hookworms) and adult liverfluke infection.

Dosage & Administration

Cattle: 2.5ml per 10kg body weight.

Sheep: 1ml per 4kg body weight.

Withdrawal periods from the last day of treatment:

Meat: 14 days | Milk: 3 days

Storage

Store in a dry place, below 30°C, protected from direct sunlight.

Keep all medicines out of reach of children.

Presentation

150ml, 500ml & 1L

FOR VETERINARY USE ONLY

KenTraz



Composition

Per ml solution, Toltrazuril 25mg

Indications

KenTraz 2.5% oral is effective against Coccidiosis in poultry and turkey.

Dosage & Administration

For oral administration in drinking water. Poultry and Turkey: 7mg per kg bodyweight per day for 2 days, 24 hours continuously or for 8 hours or each day or 100ml per 100 litres of drinking water for 2 days. Medicated drinking water should be used within 24 hours.

Withdrawal periods from last day of treatment: Poultry and Turkey 19 days. Do not use in birds producing eggs for human consumption.

Storage

Store in a dry place, below 30°C, protected from direct sunlight. Keep all medicines out of reach of children.

Presentation

100ml

FOR VETERINARY USE ONLY



KENVET AFRI LIMITED

P.o. Box 907-00606, Sarit Centre, Nairobi, Kenya.

Samovet Injection



Composition

Each Sachet of 1 gm contains:
Isometamidium Chloride Hydrochloride 1 gm

Indications

Indications, dose rates, precautions
(see enclosed direction leaflet)
For use in cattle, sheep, goats, camels, horses,
dogs and buffaloes.

Dosage & Administration

Dissolve content of 1g sachet in 50ml or 100ml
of sterile water.
Administer by deep I.M. injection.

Withdrawal Periods
Meat: 1 month
Milk: Nil

Storage

Store in a dry place, below 30°C, protected
from direct sunlight.
Keep all medicines out of reach of children.

Presentation

1gm

FOR VETERINARY USE ONLY

TetraKen Aerosol



Composition

Each TetraKen Aerosol can contains 5g
Oxytetracycline Hydrochloride
(equivalent to 3.58% w/w).

Indications

TetraKen Aerosol is indicated for treatment of:

- Foot rot in Cattle, horses, sheep, goats and pigs.
- Wound management after dehorning or surgery.
- Topical infections caused by organisms sensitive to oxytetracycline.

Dosage & Administration

- For topical application only.
- Wound to be pared and cleaned before application.
- Shake the can well before use.
- Spray for a few seconds until the wound lesion is completely covered.
- Treated animals should be allowed to stand on dry ground for one hour before returning to pasture.

No Withdrawal Periods Apply.

Storage

Store in a dry place, below 30°C, protected
from direct sunlight.
Keep all medicines out of reach of children.

Presentation

200ml

FOR VETERINARY USE ONLY



KENVET AFRI LIMITED

P.o. Box 907-00606, Sarit Centre, Nairobi, Kenya.

KenVaq Plus



Composition

Each ml contains:

Buparvaquone 50mg

Furosemide BP 55mg

Sorbitan mono oleate BP 100mg

(as preservative) Oil base

Indications

For the treatment of theileriosis (east coast fever) particularly for advanced cases with pulmonary oedema Buparvaquone kills theilerial schizonts (in lymphoid cells) and piroplasms (in red blood cells) and it suppresses preschizont stages during the incubation period of disease Buparvaquone kill the parasite through its actions on their mitochondrial electron transport (respiratory) system. Furosemide is a diuretic, which resolves pulmonary oedema.

Dosage & Administration

Injection by the Intramuscular route into the neck muscles at the rate of 1ml / 20 kg (2.5 mg per kg of buparvaquone and 2.75 mg/kg of furosemide) Repeat after 48 hours. In cases of exceptionally severe infections with pulmonary oedema, further treatment at half the dosage rate may be required at 24 hour interval.

Withdrawal Period:

Milk for human consumption should not be taken from animals treated with KenVaq Plus until atleast 48 hours after treatment.

Milk for animals treated with KenVaq Plus is safe for consumption by calves. Animals should not be slaughtered for human consumption until at least 42 days after treatment with KenVaq Plus.

Warning / Precaution:

Localised swelling may occur at Injection sites but it resolves in a few days.

KenVaq Plus is very safe so overdosage is unlikely to cause significant adverse effect. KenVaq Plus must be administered only by the Intramuscular route. Intravenous Injection may cause severe shock KenVaq Plus is poorly mobilised after subcutaneous injection and its curative effect is greatly reduced.

Furosemide is chemically similar to the sulphonamides so KenVaq Plus should not be used in animal with sulphonamide sensitivity

Storage

Store in a dry place, below 30°C, protected from direct sunlight.

Keep all medicines out of reach of children.

Presentation

50ml & 100ml

FOR VETERINARY USE ONLY



KENVET AFRI LIMITED

P.o. Box 907-00606, Sarit Centre, Nairobi, Kenya.



KENVET AFRI LIMITED

P.o. Box 907-00606, Sarit Centre, Nairobi, Kenya.